

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : A61K 38/00	A2	(11) International Publication Number: WO 00/66138 (43) International Publication Date: 9 November 2000 (09.11.00)
<p>(21) International Application Number: PCT/US00/11251</p> <p>(22) International Filing Date: 27 April 2000 (27.04.00)</p> <p>(30) Priority Data: 09/302,596 30 April 1999 (30.04.99) US</p> <p>(71) Applicant: BIONEBRASKA, INC. [US/US]; 3820 N.W. 46th Street, Lincoln, NE 68524 (US).</p> <p>(72) Inventors: COOLIDGE, Thomas, R.; 3820 N.W. 46th Street, Lincoln, NE 68524 (US). EHLERS, Mario, R., W.; 3820 N.W. 46th Street, Lincoln, NE 68524 (US).</p> <p>(74) Agent: SEASE, Edmund, J.; Zarley, McKee, Thomte, Voorhees &amp; Sease, Suite 3200, 801 Grand Avenue, Des Moines, IA 50309-2721 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>	
<p>(54) Title: METABOLIC INTERVENTION WITH GLP-1 TO IMPROVE THE FUNCTION OF ISCHEMIC AND REPERFUSED TISSUE</p> <p>(57) Abstract</p> <p>Individuals in need of treatment of ischemia-related reperfusion are treated, preferably intravenously, with a composition which includes a compound which binds to a receptor for the glucagon-like peptide-1. The invention relates to both the method and compositions for such treatment.</p>		

BEST AVAILABLE COPY

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE: METABOLIC INTERVENTION WITH GLP-1 TO IMPROVE THE  
FUNCTION OF ISCHEMIC AND REPERFUSED TISSUE

CROSS REFERENCE TO A RELATED APPLICATION

5 This application is a continuation-in-part of provisional application 60/103,498  
filed October 8, 1998.

FIELD OF THE INVENTION

This invention relates to metabolic intervention with GLP-1 to therapeutically  
10 improve the function of ischemic and reperfused tissue.

BACKGROUND OF THE INVENTION

Cellular damage to aerobic organ tissues is well recognized as a consequence of  
ischemia, whether endogenous as in the case of a spontaneous coronary artery occlusion, or  
15 iatrogenic such as with open heart, coronary bypass surgery, or transplant procedures with  
the heart or other organs such as the lung, liver, kidney, pancreas and gastrointestinal tract.  
The degree and duration of the ischemia causing events are relevant to the amount of cell  
death and/or reversible cellular dysfunction. It is also known that much of the tissue  
damage in fact occurs upon reperfusion (i.e., resumption of blood flow) and re-oxygenation  
20 of the previously anoxic tissue. Reperfusion injury has been the subject of considerable  
recent study prompted by medical advances particularly in the treatment of reperfusion  
injury after myocardial infarction and other myocardial remedial procedures such as  
coronary bypass, other open heart surgeries, as well as organ transplants.

As a side product of normal aerobic respiration, electrons are routinely lost from the  
25 mitochondrial electron transport chain. Such electrons can react with molecular oxygen to  
generate the reactive free radical superoxide which through other reaction steps in the  
presence of hydrogen peroxide and iron produces the extraordinarily reactive and toxic  
hydroxyl radical. Metabolically active aerobic tissues possess defense mechanisms  
dedicated to degrading toxic free radicals before these reactive oxygen species can interact  
30 with cellular organelles, enzymes, or DNA, the consequences of which could, without such  
protective mechanisms, be cell death. These defense mechanisms include the enzymes

superoxide dismutase (SOD) which disproportionates superoxide, catalase which degrades hydrogen peroxide, and the peptide glutathione which is a non-specific free radical scavenger.

While not fully understood, it is believed that with ischemia of metabolic tissues  
5 and subsequent reperfusion, a complex group of events occurs. Initially during the  
ischemic period, intracellular anti-oxidant enzyme activity appears to decrease, including  
that of SOD, catalase, and glutathione. There is also an indication that the level of  
xanthine oxidase activity concomitantly increases in vascular endothelial tissue during the  
ischemic event. The combination of enhanced ability to produce oxygen free radicals (via  
10 enhanced xanthine oxidase activity) and reduced ability to scavenge the same oxygen  
radicals (via reduced SOD, catalase and glutathione activity) greatly sensitizes the ischemic  
cell to an oxidative burst, and hence damage, should these cells be subsequently reperfused  
with blood and therefore oxygen. This oxidative burst occurring within seconds to minutes  
of reperfusion could result in reversible and irreversible damage to endothelial cells and  
15 other cells constituting the ischemic-reperfused organ matrix. If, for example, the heart is  
the organ under consideration, reversible oxidative damage can contribute to myocardial  
stunning, whereas irreversible damage presents itself as a myocardial infarction. Attendant  
with this initial oxidative burst is oxidation damage to cell membranes. Lipid oxidation in  
cell membranes appears to play a role in neutrophil chemotaxis to post-ischemic areas.  
20 Such activated neutrophils adhere to vascular endothelium, induce the conversion of  
xanthine dehydrogenase to xanthine oxidase within said endothelial cells, and further  
aggravate loss of endothelial integrity. Activated neutrophils also migrate out of the  
vasculature into myocardial interstitial spaces where the inflammatory cells can directly  
kill myocytes. Additionally, perturbations in normal calcium mobilization from  
25 sarcoplasmic reticulum as a consequence of ischemia-reperfusion contribute to reversible  
myocardial dysfunction referred to as myocardial stunning.

The consequences of ischemia-reperfusion events are reversible and irreversible  
cell damage, cell death, and decreased organ functional efficiency. More specifically, in  
the case of myocardial reperfusion injury, the consequences include myocardial stunning,  
30 arrhythmias, and infarction, and as a result, cardiogenic shock and potentially congestive  
heart failure.

The paradox of cellular damage associated with a limited period of ischemic anoxia followed by reperfusion is that cell damage and death appear not only likely to directly result from the period of oxygen deprivation but, additionally, as a consequence of re-oxygenation of tissues rendered highly sensitive to oxidative damage during the ischemic  
5 period. Reperfusion damage begins with the initial oxidative burst immediately upon reflow and continues to worsen over a number of hours as inflammatory processes develop in the same post-ischemic tissues. Efforts dedicated to decreasing sensitivity of post-anoxic cells to oxidative damage and, additionally, efforts to reduce inflammatory responses in these same tissues have been shown to reduce the reversible and irreversible  
10 damage to post-anoxic reperfused organs. A combination of methods to reduce both the initial oxidative burst and subsequent inflammation associated damage could provide synergistic protection against reperfusion injury.

With respect to the treatment of ischemia coincident with MI patients, common therapies now used are to employ thrombolytics such as streptokinase and t-PA and  
15 angioplasty. U.S. Pat. No. 4,976,959 discloses the administration of t-PA and SOD to inhibit tissue damage during reperfusion and/or percutaneous transluminal coronary angioplasty coincident with ischemia to restore regional blood flow. Thus, an increasing number of patients are being exposed to the likelihood of reperfusion injury and its effects, particularly cardiac patients.

20 Reperfusion injury to organs other than the heart will generally manifest itself in substantially reduced efficiency of function, a consequence of which may be premature degeneration of the organ, or simply shutdown. Additionally, transplanted organs experience enhanced rejection rates if there is significant underlying reperfusion injury.

As discussed briefly above, while the precise mechanism of reperfusion injury has  
25 not been clearly defined, mounting data, most of which has been gathered in various cardiac model studies, indicate that the generation of oxygen-derived free radicals, including superoxide anion ( $O_2^-$ ), the hydroxyl free radical ( $\cdot OH$ ) and  $H_2O_2$ , results as a consequence of the reintroduction of molecular oxygen with reperfusion and plays an important role in tissue necrosis. Agents which either decrease the production of these  
30 oxygen derived free radicals (including allopurinol and deferoxamine) or increase the degradation of these materials such as superoxide dismutase, catalase, glutathione, and

copper complexes, appear to limit infarct size and also may enhance recovery of left ventricular function from cardiac stunning.

The use of metabolic intervention as a therapy specifically during acute myocardial infarction is well established, although not without controversy. There is abundant  
5 experimental and clinical evidence to support the use of a glucose-insulin-potassium (GIK) infusion - the primary form of metabolic intervention - after acute MI, particularly following the success of the Swedish DIGAMI study (Malmberg, K, and DIGAMI Study Group (1997) Prospective randomized study of intensive insulin treatment on long term  
10 survival after acute myocardial infarction in patients with diabetes mellitus. *Brit. Med. J.* 314, 1512-1515). The DIGAMI study emphasized the efficacy of a glucose-insulin infusion for acute MI in diabetic patients, but this type of therapy has never been suggested or used for reperfusion.

It therefore can be seen that there is a need for a safe effective composition having broad applicability to prevent or ameliorate the harmful effects of ischemia and reperfusion  
15 for tissues in general, especially organ tissue and, including but not limited to myocardium. It is primary object of the present invention to fulfill this need.

Another object of the present invention is to provide a method for treating ischemia and reperfusion without the side effects normally attendant with therapies presently available.

20 Still another object of the present invention is to provide a pharmaceutically acceptable carrier composition which can be used for intravenous administration of the compositions of the present invention without any significant undesirable side effects and without adversely affecting antigenic or immune stimulating properties.

These and other objects and benefits of the present invention will be apparent to  
25 those skilled in the art from the further description and the accompanying claims.

#### SUMMARY OF THE INVENTION

Individuals in need of treatment of ischemia and/or reperfusion are treated, preferably intravenously, with a composition which includes a compound which binds to a  
30 receptor for the glucagon-like peptide-1. The invention relates to both the method and compositions for such treatment.

DETAILED DESCRIPTION OF THE INVENTION

GLP-1 is a glucose-dependent insulinotropic hormone that effectively enhances peripheral glucose uptake without inducing dangerous hypoglycemia. Further, GLP-1 strongly suppresses glucagon secretion, independent of its insulinotropic action, and thereby powerfully reduces plasma free fatty acid (FFA) levels substantially more than can be accomplished with insulin. High FFA levels have been implicated as a major toxic mechanism during myocardial ischemia.

We have now developed the concept of GLP-1 as a metabolic therapy for ischemia-reperfusion injury. This development was based on the realization that there are two clinical situations in which ischemia-reperfusion is a routine, and potentially dangerous, event: thrombolytic procedures for acute MI, and cardiac reperfusion following ischemic cardioplegia during heart surgery. Moreover, recent experimental and clinical data have established that the phenomenon of ischemia-reperfusion is particularly responsive to metabolic therapy with GIK infusion, even more so than isolated ischemia without reperfusion (Apstein, CS (1998) Glucose-insulin-potassium for acute myocardial infarction. Remarkable results from a new prospective, randomized trial. *Circulation* 98, 2223-2226).

The two most important therapeutic advances in the treatment of acute ischemia coincident with MI in the past decade have been the introduction of thrombolysis and  $\beta$ -blockade. However, despite this overall success, some studies of thrombolysis have revealed an early excess mortality, which has been attributed to reperfusion-induced injury and myocardial stunning. The mechanisms underlying stunning are complex, but an emerging consensus is that this is likely related to intracellular acidosis leading to dysfunctional sarcolemmal  $Ca^{2+}$  pumps and cytosolic  $Ca^{2+}$  overload. The net result is impaired myocardial contractile function leading to decreased mechanical efficiency, as well as reperfusion ventricular arrhythmias. Moreover, recent research has established that the intracellular acidosis, in turn, is due to an imbalance between glycolysis and complete glucose oxidation, in the sense that the rate of glycolysis is uncoupled from the oxidation of pyruvate (the end product of glycolysis) in the TCA cycle. This uncoupling results in net  $H^+$  production due to conversion of pyruvate to lactate. The most likely cause for this imbalance is the presence of high plasma free fatty acid (FFA) levels, which preferentially

enter the mitochondria and inhibit pyruvate oxidation, a mechanism that elegantly accounts for the well-established observation that hearts perfused with FFA are less able to recover in the reperfusion phase than hearts perfused with glucose. It has here been discovered, and is one of the bases of this therapeutic invention that GLP-1 suppresses FFA beyond  
5 what is expected with insulin which is at the 50% level of suppression, and GLP-1 can be as high as 90% suppression of FFA.

These considerations have strengthened our conviction to treat ischemia-reperfusion with glucagon-like peptides. It is well established that during normal perfusion and adequate oxygenation, the heart depends on aerobic metabolism and uses FFAs as its  
10 preferred fuel. In contrast, during ischemia (reduced blood flow) or hypoxia (reduced O<sub>2</sub> tension),  $\beta$ -oxidation of fatty acids is impaired (because it is strictly aerobic) and continued provision of ATP is dependent increasingly on anaerobic glycolysis. During the ischemic period, glucose-insulin is of benefit because it enhances glucose uptake and stimulates glycolysis, thereby providing ATP for maintenance of essential membrane functions,  
15 especially ion transport. Moreover, glucose-insulin suppresses adipose tissue lipolysis, thereby reducing plasma FFA levels and uptake of FFAs into the myocardium. High levels of FFAs are toxic to the ischemic myocardium, both by direct detergent effects on membranes and increases in cAMP, and by accumulation of acylcarnitine, which inhibits Ca<sup>2+</sup> pumps. The net effect is disturbance of ion exchange, cytosolic Ca<sup>2+</sup> overload, and  
20 resultant contractile dysfunction and arrhythmias.

During the reperfusion period, glucose-insulin is of benefit because, as explained above, this therapy can alleviate the metabolic imbalance that produces stunning. This is achieved by direct stimulation of PDH and hence pyruvate oxidation, and indirectly by  
reduced FFA uptake and hence improved ratio of pyruvate to FFA oxidation.

25 From the above discussion it is evident that the dual action of glucose-insulin - enhanced glucose uptake and metabolism, and reduced FFA levels - has substantial therapeutic potential in reperfusion. Some have expressed a concern that during profound, essentially zero-flow ischemia, glycolytic end products, namely lactate, will accumulate due to inadequate "wash-out". Lactate accumulation, in turn, leads to high intracellular  
30 proton concentrations, and failure to reoxidize NADH; high [H<sup>+</sup>] and NADH/NAD<sup>+</sup> ratios inhibit productive glycolysis. Under these circumstances, glucose can be toxic to cells,



because ATP is actually consumed in the production of fructose-1,6-bisphosphate, and high  $[H^+]$  can aggravate myocyte necrosis (Neely, JR, and Morgan, HE (1974) Relationship between carbohydrate and lipid metabolism and the energy balance of heart muscle. *Ann. Rev. Physiol.* 36, 413-459). However, these concerns have not been borne out by the weight of experimental and clinical data, which indicate that glucose-insulin produces beneficial results. While not wishing to be bound by theory, the likely explanation for this is that in humans, acute spontaneous ischemia is not a condition of zero-flow ischemia, but instead represents a region of low-flow ischemia in which residual perfusion is adequate for substrate delivery and lactate washout. This realization has now provided a powerful physiological logic for the use of metabolic therapy in ischemia-reperfusion.

Modern cardiac surgery, whether involving cardiac valve replacement or coronary artery bypass grafting (CABG), routinely requires hypothermic cardioplegic arrest, aortic crossclamping, and cardiopulmonary bypass during surgery. Effectively, therefore, routine cardiac surgery induces a state of elective global ischemia followed by reperfusion, which potentially exposes the heart to all the attendant risks and injuries peculiar to myocardial ischemia-reperfusion. Hence, prevention of myocardial damage during and after cardiac operations remains a major concern. Elective cardioplegic ischemia followed by reperfusion has obvious parallels with ischemia-reperfusion encountered during acute MI followed by revascularization, and thus many of the pathophysiological principles considered in previous sections also apply during cardiac surgery. However, there are some notable differences between surgical cardioplegic ischemia-reperfusion and MI-associated ischemia-reperfusion. During surgery, the heart is arrested (cardioplegia) and infused with a cold (hypothermic) solution designed to optimize myocardial preservation. After completion of the surgery, the heart is reactivated and reperfused with oxygenated blood at body temperature. This produces a sequence of hypothermic ischemia and normothermic reperfusion, which may prevent the accumulation of high tissue levels of  $H^+$  and lactate. Moreover, unlike acute MI, hypothermic cardioplegia represents a state of global, zero-flow ischemia, followed by global reperfusion.

In our previous application (Serial No. 60/103,498), of which this is a continuation-in-part, we reviewed the disadvantages of glucose-insulin infusions and the advantages of substituting these with a GLP-1 infusion, which is safer than insulin. In summary, GIK

infusions carry significant risks of both hypoglycemia and hyperglycemia, and are technically demanding and staff-intensive. The dangers of hypoglycemia are obvious.

In contrast, these risks do not exist with a GLP-1 infusion. Glucagon-like peptide (7-36) amide (GLP-1) is a natural, gut-derived, insulinotropic peptide that constitutes a major component of the so-called incretin effect. GLP-1 exerts its major effect at the pancreatic endocrine cells, where it (1) regulates insulin expression and secretion from the  $\beta$ -cells in a glucose-dependent fashion; (2) stimulates the secretion of somatostatin; and (3) suppresses the secretion of glucagon from the  $\alpha$  cells. Although not formally resolved, the strong glucagonostatic effect is presumed to result from one or all of the following: (1) direct suppression by stimulation of GLP-1 receptors on  $\alpha$  cells, although this is unlikely; (2) paracrine suppression of glucagon secretion by intra-islet release of somatostatin; or (3) paracrine suppression by intra-islet release of insulin. Whatever the cellular mechanism, GLP-1 is unique in its capacity to simultaneously stimulate insulin secretion and inhibit glucagon release. Although a therapeutic insulin infusion also inhibits glucagon release, this effect is not as potent as that of GLP-1, which exerts a direct, intra-islet paracrine inhibition of glucagon secretion.

The dual capacity of GLP-1 to powerfully stimulate insulin release and inhibit glucagon secretion, together with the strict glucose-dependence of its insulinotropic action, endow this molecule with a unique therapeutic potential in the management of ischemia-reperfusion. First, GLP-1 strongly stimulates the secretion of endogenous insulin and therefore can be used to achieve all of the beneficial actions attributed to an insulin infusion in the metabolic treatment of ischemia-reperfusion. Although high-dose GIK infusions typically contain 25-33% glucose and 50-100 U insulin/L, the requirement for introduction of hyperglycemia per se to achieve therapeutic efficacy, versus only providing a metabolic milieu for the safe administration of high doses of insulin, is unclear. It is likely that adequate blood glucose levels are required to enable substrate delivery, but this does not necessarily imply a need for hyperglycemia and should not detract from the fact that insulin exerts important effects other than glucose uptake. Therefore, a therapeutic GLP-1 infusion will likely only require a modest (e.g., 5%) glucose coinfusion in order to maintain blood glucose at slightly above physiological levels in order to trigger insulin release. Glucose is not required as a safety measure, since blood levels of  $\leq 3.5$  mM

abrogate the insulin-stimulating activity of GLP-1, thereby completely protecting against the dangers of hypoglycemia.

Second, GLP-1 exerts a powerful glucagonostatic effect, which together with its insulinotropic action will lead to a strong suppression of FFAs. One of the major benefits of glucose-insulin infusions is the reduction in circulating FFA levels and the suppression of FFA uptake. FFAs and their metabolites have direct toxic effects on the ischemic myocardium as well as during the reperfusion period, when they contribute to stunning, and hence reduction of FFA levels is a major therapeutic goal of metabolic intervention in ischemia-reperfusion, goal of metabolic intervention in ischemia-reperfusion. As glucagon is a powerful stimulus for adipose tissue lipolysis and FFA production, GLP-1 mediated glucagon suppression further augments the insulin-induced reduction in circulating FFAs. Thus, GLP-1 therapy is superior to a glucose-insulin infusion in this regard. Indeed, preliminary data in healthy volunteers indicate that an intravenous GLP-1 infusion will reduce fasting plasma FFA levels to <10% of control values.

GLP-1 should be effective in the majority of patients without requiring concurrent glucose administration. However, a small proportion of subjects may require glucose/GLP-1 to elicit an adequate insulin response. In addition, it also may be necessary to administer potassium to correct excess shifts of potassium in the intracellular compartment when glucose is co-administered with GLP-1.

In addition to GLP-1 or its biological analogues, the therapy can include use of free radical scavengers such as glutathione, melatonin, Vitamin E, and superoxide dismutase (SOD). In such combinations reperfusion damage risk is lessened even further.

The term "GLP-1", or glucagon-like peptide, includes mimetics, and as used in the context of the present invention can be comprised of glucagon-like peptides and related peptides and analogs of glucagon-like peptide-1 that bind to a glucagon-like peptide-1 (GLP-1) receptor protein such as the GLP-1 (7-36) amide receptor protein and has a corresponding biological effect on insulin secretion as GLP-1 (7-36) amide, which is a native, biologically active form of GLP-1. See Goke, B and Byrne, M, *Diabetic Medicine*, 1996, 13:854-860. The GLP-1 receptors are cell-surface proteins found, for example, on insulin-producing pancreatic  $\beta$ -cells. Glucagon-like peptides and analogs will include species having insulinotropic activity and that are agonists of, *i.e.* activate, the GLP-1

receptor molecule and its second messenger activity on, *inter alia*, insulin producing  $\beta$ -cells. Agonists of glucagon-like peptide that exhibit activity through this receptor have been described: EP 0708179A2; Hjorth, S.A. *et al.*, *J. Biol. Chem.* 269 (48):30121-30124 (1994); Siegel, E.G. *et al.* Amer. Diabetes Assoc. 57th Scientific Sessions, Boston (1997);  
5 Hareter, A. *et al.* Amer. Diabetes Assoc. 57th Scientific Sessions, Boston (1997); Adelhorst, K. *et al.* *J. Biol. Chem.* 269(9):6275-6278 (1994); Deacon C.F. *et al.* 16th International Diabetes Federation Congress Abstracts, *Diabetologia Supplement* (1997); Irwin, D.M. *et al.*, *Proc. Natl. Acad. Sci. USA.* 94:7915-7920 (1997); Mosjov, S., *Int. J. Peptide Protein Res.* 40:333-343 (1992). Glucagon-like molecules include polynucleotides  
10 that express agonists of GLP-1, i.e. activators of the GLP-1 receptor molecule and its secondary messenger activity found on, *inter alia*, insulin-producing  $\beta$ -cells. GLP-1 mimetics that also are agonists include, for example, chemical compounds specifically designed to activate the GLP-1 receptor. Glucagon-like peptide-1 antagonists are also known, for example see e.g. Watanabe, Y *et al.*, *J. Endocrinol.* 140(1):45-52 (1994), and  
15 include exendin (9-39) amine, an exendin analog, which is a potent antagonist of GLP-1 receptors (see, e.g. WO97/46584). Recent publications disclose Black Widow GLP-1 and Ser<sup>2</sup> GLP-1, see G.G. Holz, J.F. Hakner/*Comparative Biochemistry and Physiology*, Part B 121(1998)177-184 and Ritzel, *et al.*, *A synthetic glucagon-like peptide-1 analog with improved plasma stability*, *J. Endocrinol* 1998 Oct. 159(1):93-102.

20 Further embodiments include chemically synthesized glucagon-like polypeptides as well as any polypeptides or fragments thereof which are substantially homologous. "Substantially homologous," which can refer both to nucleic acid and amino acid sequences, means that a particular subject sequence, for example, a mutant sequence, varies from a reference sequence by one or more substitutions, deletions, or additions, the net  
25 effect of which does not result in an adverse functional dissimilarity between reference and subject sequences. For purposes of the present invention, sequences having greater than 50 percent homology, and preferably greater than 90 percent homology, equivalent biological activity in enhancing  $\beta$ -cell responses to plasma glucose levels, and equivalent expression characteristics are considered substantially homologous. For purposes of determining  
30 homology, truncation of the mature sequence should be disregarded. Sequences having

lesser degrees of homology, comparable bioactivity, and equivalent expression characteristics are considered equivalents.

Mammalian GLP peptides and glucagon are encoded by the same gene. In the ileum the phenotype is processed into two major classes of GLP peptide hormones, namely  
5 GLP-1 and GLP-2. There are four GLP-1 related peptides known which are processed from the phenotypic peptides. GLP-1 (1-37) has the sequence His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ. ID NO:1). GLP-1 (1-37) is amidated by post-translational processing to yield GLP-1 (1-36) NH<sub>2</sub>, which has the sequence His Asp  
10 Glu Phe Glu Arg His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH<sub>2</sub>) (SEQ. ID NO:2); or is enzymatically processed to yield GLP-1 (7-37) which has the sequence His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ. ID NO:3). GLP-1 (7-37) can also be amidated to yield  
15 GLP-1 (7-36) amide which is the natural form of the GLP-1 molecule, and which has the sequence His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH<sub>2</sub>) (SEQ. ID NO:4) and in the natural form of the GLP-1 molecule.

Intestinal L cells secrete GLP-1 (7-37) (SEQ. ID NO:3) and GLP-1 (7-36) NH<sub>2</sub>  
20 (SEQ. ID NO:4) in a ratio of 1 to 5, respectively. These truncated forms of GLP-1 have short half-lives *in situ*, i.e., less than 10 minutes, and are inactivated by an aminodipeptidase IV to yield Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ. ID NO:5); and Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala  
25 Trp Leu Val Lys Gly Arg (NH<sub>2</sub>) (SEQ. ID NO:6), respectively. The peptides Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ. ID NO:5) and Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH<sub>2</sub>) (SEQ. ID NO:6), have been speculated to affect hepatic glucose production, but do not stimulate the  
30 production or release of insulin from the pancreas.

There are six peptides in Gila monster venoms that are homologous to GLP-1. Their sequences are compared to the sequence of GLP-1 in Table 1.

TABLE 1

- a. HAEGTFTSDVSSYLEGQAAKEFIAWLVKGRNH<sub>2</sub>
- b. HSDGTFTSDLSKQMEEEAVRLFIEWLKNNGPSSGAPPPSNH<sub>2</sub>
- 5 c. DLSKQMEEEAVRLFIEWLKNNGPSSGAPPPSNH<sub>2</sub>
- d. HGEFTFTSDLSKQMEEEAVRLFIEWLKNNGPSSGAPPPSNH<sub>2</sub>
- e. HSDATFTAAYSKLLAKLALQKYLELGSSTSPRPPS
- f. HSDATFTAAYSKLLAKLALQKYLELGSSTSPRPPS
- g. HSDAIFTTEYSKLLAKLALQKYLASILGSRTSPPPNH<sub>2</sub>
- 10 h. HSDAIFTQQYSKLLAKLALQKYLASILGSRTSPPPNH<sub>2</sub>

- a=GLP-1(SEQ. ID NO:4).
- b=Exendin 3(SEQ. ID NO:7).
- c=Exendin 4(9-39)(NH<sub>2</sub>(SEQ.ID NO:8).
- 15 d=Exendin 4(SEQ.ID NO:9).
- e=Helospectin I(SEQ.ID NO:10).
- f=Helospectin II(SEQ. ID NO:11).
- g=Helodermin(SEQ. ID NO:12).
- h=Q<sup>8</sup>, Q<sup>9</sup> Helodermin(SEQ. ID No:13).

The major homologies as indicated by the outlined areas in Table 1 are: peptides c and h are derived from b and g, respectively. All 6 naturally occurring peptides (a, b, d, e, f and g) are homologous in positions 1, 7, 11 and 18. GLP-1 and exendins 3 and 4 (a, b and d) are further homologous in positions 4, 5, 6, 8, 9, 15, 22, 23, 25, 26 and 29. In position 2, A, S and G are structurally similar. In position 3, residues D and E (Asp and Glu) are structurally similar. In positions 22 and 23 F (Phe) and I (Ile) are structurally similar to Y (Tyr) and L (Leu), respectively. Likewise, in position 26 L and I are structurally equivalent.

Thus, of the 30 residues of GLP-1, exendins 3 and 4 are identical in 15 positions and equivalent in 5 additional positions. The only positions where radical structural changes are evident are at residues 16, 17, 19, 21, 24, 27, 28 and 30. Exendins also have 9 extra residues at the carboxyl terminus.

The GLP-1 like peptides can be made by solid state chemical peptide synthesis. GLP-1 can also be made by conventional recombinant techniques using standard procedures described in, for example, Sambrook and Maniatis. "Recombinant," as used herein, means that a protein is derived from recombinant (e.g., microbial or mammalian) expression systems which can be genetically modified to contain an expression gene for GLP-1 or its biologically active analogues.

The GLP-1 like peptides can be recovered and purified from recombinant cell cultures by methods including, but not limited to, ammonium sulfate or ethanol) precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High performance liquid chromatography (HPLC) can be employed for final purification steps.

The polypeptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from prokaryotic or eukaryotic hosts (for example by bacteria, yeast, higher plant, insect and mammalian cells in culture or *in vivo*). Depending on the host employed in a recombinant production procedure, the polypeptides of the present invention are generally non-glycosylated, but may be glycosylated.



GLP-1 activity can be determined by standard methods, in general, by receptor-binding activity screening procedures which involve providing appropriate cells that express the GLP-1 receptor on their surface, for example, insulinoma cell lines such as RINmSF cells or INS-1 cells. See also Mosjov, S.(1992) and EP0708170A2. In addition  
5 to measuring specific binding of tracer to membrane using radioimmunoassay methods, cAMP activity or glucose dependent insulin production can also be measured. In one method, a polynucleotide encoding the receptor of the present invention is employed to transfect cells to thereby express the GLP-1 receptor protein. Thus, for example, these methods may be employed for screening for a receptor agonist by contacting such cells  
10 with compounds to be screened and determining whether such compounds generate a signal, i.e. activate the receptor.

Polyclonal and monoclonal antibodies can be utilized to detect, purify and identify GLP-1 like peptides for use in the methods described herein. Antibodies such as ABGA1178 detect intact unspliced GLP-1 (1-37) or N-terminally-truncated GLP-1 (7-37)  
15 or (7-36) amide. Other antibodies detect on the very end of the C-terminus of the precursor molecule, a procedure which allows by subtraction to calculate the amount of biologically active truncated peptide, i.e. GLP-1 (7-37) or (7-36) amide (Orskov et al. Diabetes, 1993, 42:658-661; Orskov et al. J. Clin. Invest. 1991, 87:415-423).

Other screening techniques include the use of cells which express the GLP-1  
20 receptor, for example, transfected CHO cells, in a system which measures extracellular pH or ionic changes caused by receptor activation. For example, potential agonists may be contacted with a cell which expresses the GLP-1 protein receptor and a second messenger response, e.g. signal transduction or ionic or pH changes, may be measured to determine whether the potential agonist is effective.

25 The glucagon-like peptide-1 receptor binding proteins of the present invention may be used in combination with a suitable pharmaceutical carrier. Such compositions comprise a therapeutically effective amount of the polypeptide, and a pharmaceutically acceptable carrier or excipient. Such a carrier includes, but is not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, lactose, phosphate, mannitol, arginine,  
30 trehalose and combinations thereof. The formulations should suit the mode of administration and are readily ascertained by those of skill in the art. The GLP-1 peptide

may also be used in combination with agents known in the art that enhance the half-life *in vivo* of the peptide in order to enhance or prolong the biological activity of the peptide. For example, a molecule or chemical moiety may be covalently linked to the composition of the present invention before administration thereof. Alternatively, the enhancing agent  
5 may be administered concurrently with the composition. Still further, the agent may comprise a molecule that is known to inhibit the enzymatic degradation of GLP-1 like peptides may be administered concurrently with or after administration of the GLP-1 peptide composition. Such a molecule may be administered, for example, orally or by injection.

10 Patients administered GLP-1 or its analogues in combination with the carrier systems here enumerated, especially those treated before a planned event or within the first 4 hours after an ischemic event, are observed to have less arrhythmia, less tissue damage, and less discomfort without side effects.

From these considerations it is evident that an infusion of GLP-1 can be expected to  
15 exert a major therapeutic effect in myocardial reperfusion. It is expected that GLP-1 can be administered either by I.V. or subcutaneous administration for continuous infusion by intravenous (I.V.) 0.1 pmol/kg/min to 10 pmol/kg/min and by subcutaneous (S.C.) 0.1 pmol/kg/min to 75 pmol/kg/min, and for single injection (bolus) by I.V. 0.005 nmol/kg to 20 nmol/kg and S.C. 0.1 nmol/kg to 100 nmol/kg are suitable levels of administration. The  
20 GLP-1 infusion can be coadministered with glucose (5%) if required to maintain blood glucose levels  $\geq 5$  mM (to maintain efficient insulin secretion). Similarly, coadministration of potassium ( $K^+$ ) will also be considered, depending on the extent to which activation of the membrane  $Na^+/K^+$  ATPase leads to a shift of  $K^+$  into the intracellular space. The GLP-1 treatment will be commenced as early in the post-ischemic  
25 period as possible after, for example, acute spontaneous ischemia in the home or ambulance context and before reperfusion therapies, and continued thereafter. In the case of cardiac surgery, the GLP-1 infusion should commence 12-24 hours prior to surgery, during surgery from the onset of anesthesia until aortic crossclamping, and immediately after unclamping for a period of at least 72 hours postoperatively. As earlier explained, co-  
30 administration of a free radical scavenger will further aid reperfusion recovery.

From the above it can be seen that the invention accomplishes all of its stated objectives.

What is claimed is:

1. A method for amelioration of organ tissue injury caused by reperfusion of blood flow following a period of ischemia, which comprises: administering to an individual in  
5 need of such treatment an effective amount of a composition which includes a compound which binds to a receptor for glucagon-like peptide-1, in a pharmaceutical carrier.
2. The method of claim 1 wherein the glucagon-like peptide-1 is GLP-1 or a biologically active analogue thereof.
- 10 3. The method of claim 1 wherein the pharmaceutical carrier is selected from the group consisting of saline, buffered saline, dextrose, water, glycerol, ethanol, lactose, phosphate, mannitol, arginine, trehalose, and combinations thereof.
- 15 4. The method of claim 1 wherein the administering to an individual in need of treatment is at a dose level of 0.1 pmol/kg/min. up to 10 pmol/kg/min.
5. The method of claim 4 wherein there is concurrent administration of glucose.
- 20 6. The method of claim 5 wherein there is concurrent administration of potassium.
7. The method of claim 4 wherein there is concurrent administration of a free radical scavenger.
- 25 8. The method of claim 1 wherein administration commences within 4 hours of an ischemic event.
9. The method of claim 8 wherein administration occurs within 4 hours of an ischemic event and continues thereafter.
- 30 10. The method of claim 1 wherein administration is intravenously.

11. The method of claim 1 wherein administration is by subcutaneous or micropressure injection, deep lung insufflation, external or implant pump, depot injection, and other sustained release mechanisms, oral delivery and patch, buccal and other cross skin and membrane mechanisms.
- 5
12. The method of claim 1 wherein the organ tissue is the myocardium.
13. The method of claim 8 wherein administration occurs as soon as possible after the event.
- 10
14. The method of metabolic intervention with GLP-1 to improve the function of ischemic and reperfused tissue, said method comprising: administering to an individual in need of such treatment an effective amount of a composition comprising GLP-1 in a pharmaceutical carrier.
- 15
15. The method of claim 14 wherein the glucagon-like peptide-1 is GLP-1 or a biologically active analog thereof.
16. The method of claim 14 wherein the pharmaceutical carrier is selected from the group consisting of saline, buffered saline, dextrose, water, glycerol, ethanol, lactose, phosphate, mannitol, arginine, trehalose, and combinations thereof.
- 20
17. The method of claim 14 wherein the administering to an individual in need of treatment is at a dose level of 0.1 pmol/kg/min to 10 pmol/kg/min.
- 25
18. The method of claim 17 wherein there is concurrent administration of glucose.
19. The method of claim 14 wherein administration commences within 4 hours of an ischemic event.
- 30

20. The method of claim 19 wherein administration occurs within 4 hours of an ischemic event and continues thereafter.
21. The method of claim 14 wherein the need for amelioration of tissue damage by metabolic intervention arises from a medical procedure that is a surgical event selected from the group consisting of cardiac surgical procedures, organ transplants, traumatic limb amputation and reattachment.
22. The method of claim 14 wherein the medical procedure involves an ischemic reperfusion event, said event being concurrent with gut infarct and myocardial infarct.
23. A composition for use in metabolic intervention with GLP-1 to improve the function of ischemic and reperfused tissue, comprising: an effective amount of GLP-1 in combination with a pharmaceutically effective carrier.

15